

---

## Pleiades Dances Iii Iv V Ongaku

**program of special performance - yasuko-fukuda** - edvard hagerup grieg (1843-1907): "little bird" from lyric pieces book iii op.43 (1888) ... (1953-): "interlude with birds" from pleiades dances ii op. 28 (1987) program of special performance. 18 1. program (1) haru-no umi (2) godan ginuta (3) performance explaining the depiction of natural elements through sound a. koto and shamisen b. shakuhachi (4) koto solo: isuzugawa (5) shakuhachi ... **copie de femalerituals - universit  de montr al** - ceremonial was originally determined by observation of the pleiades and the moon (tooker 1970:147) and associated with the return from the fall hunt, and in the last two centuries has come to be associated with the winter solstice. it is now generally held in january. women sponsor one of the most important of the men's medicine societies, the false face ritual (kurath 1976:123), in which ... **john fonville's microtonal flutes - musicweb.ucsd** - striae ii & iii reiko manabe dustin donahue music for sarah featuring excerpts from 16 walking dances choreographed and performed by sarah brumgart 1. lisa cella 2. anne la berge 3. michael matsuno 4. rachel beetz 5. jane rigler 6. elizabeth mc nutt. songs of sappho the lyric poem and now i begin [ptolemy diatonic malakon] come holy tortoise shell my lyre and become a poem. i begin with words ... **chemical tagging of clusters in the galah survey: two ...** - mnras 000,1{9(2017) preprint 7 march 2017 compiled using mnras latex style le v3.0 chemical tagging of clusters in the galah survey: two populations and new members in the pleiades **arxiv:0707.3144v2 [astro-ph] 26 nov 2007** - (an et al. 2007, hereafter paper iii). we extend the hyades-based empirical corrections on the- oretical isochrones using the pleiades and apply our tech- niques to a set of cepheid-bearing open clusters that have good multi-color photometry. selection of cluster sample and compilation of cluster photometry are presented in § 2. proce- dures on the isochrone calibration and ms-fitting method are ... -<sup>1</sup> **model atmosphere abundance analysis** - model atmosphere abundance analysis of eight stars in the pleiades peter s. conti and stephen e. strom august 1967 ... iii. choice of model parameters in order to perform a model atmosphere analysis of these stars, it is necessary to obtain trial values of the parameters specifying the model, namely, the effective temperature  $t$  and the surface gravity  $g$ . ef f